

THE COMMONWEALTH OF MASSACHUSETTS

EXECUTIVE OFFICE OF ENERGY AND ENVIRONMENTAL AFFAIRS



Department of Agricultural Resources

251 Causeway Street, Suite 500, Boston, MA 02114
617-626-1700 fax: 617-626-1850 www.mass.gov/agr



THE ACT PROTECTING CHILDREN AND FAMILIES FROM HARMFUL PESTICIDES OF 2000

Massachusetts Pesticide Enforcement Consumer Information Bulletin FOR SCHOOLS, DAYCARE CENTERS AND SCHOOL AGE CHILD CARE PROGRAMS

The Massachusetts Pesticide Control Act requires parents, staff, and children to receive this Consumer Information bulletin whenever pesticide applications are being made on the property of your school, daycare center or school age child care program. This bulletin is being provided to you along with a Standard Written Notification form and a Pesticide Specific Factsheet.

Why am I receiving this information and what should I do when I receive it?

The purpose of the Standard Written Notification is to provide you with information about pesticide applications which are taking place on the property of your school, day care center or school age child care program. The bulletin provides information about precautions you can take to minimize exposure to any pesticides. The Pesticide Specific Factsheets provide information about the properties of the pesticides being used.

Who applies pesticides in my school, daycare center or school age child care program?

Commercial pest management professionals, facilities managers, grounds personnel or custodians. Regardless of the approach used, the person who applies the pesticides must have a current and valid Pesticide Bureau Applicator license. Check the standard written notification form for the applicator's license number.

How do I know when pesticides are being applied?

Employees, supervised children and their guardians must receive standard written notification at least two working days prior to the application of pesticides outdoors on the property. The standard written notification form, which accompanies this bulletin, includes:

- approximate dates when the application shall commence and conclude;
- specific location of the application;
- product name, type and EPA Registration number of the pesticide;
- a Pesticide Specific Fact Sheet;
- a description of the purpose of the application and
- this Consumer Information Bulletin

The notification must also be posted in a common area of the facility at least two working days before the outdoor application is to commence and at least 72 hours after the application. Treated areas will be posted with clear and conspicuous warning signs along the perimeter. This information will be supplied to the school by the licensed pesticide applicator.

Are applications of pesticides safe?

All pesticides must be treated with caution. They are intended to be specifically poisonous to target pest insects, weeds, mold, fungus etc. - and may also be harmful to other living things including humans. Some degree of risk is always posed by their use. Because of this inherent risk, a number of regulatory and non-regulatory mechanisms have evolved to deal with those risks. Included among these mechanisms are pesticide regulations such as those enforced by Massachusetts Pesticide Enforcement; licensing and training of pesticide applicators; improved pesticide application methods; and the use of Integrated Pest Management (IPM).

What precautions can I take to minimize my exposure to pesticide applications?

There are several precautions that can be taken to reduce potential exposure to pesticides. These precautions will vary depending on where and how the pesticides are applied. Chemicals may be ingested, inhaled and absorbed through the skin. Know where the pesticide will be applied and how you might come into contact with it. Use common sense. The licensed pesticide applicator is required to post yellow signs to indicate a pesticide application on school grounds. These are some suggested general precautions. Ask the licensed pesticide applicator for other suggestions or directions specific to the work being done.

For outdoor applications:

- be familiar with the small yellow signs which applicators are required to post when a pesticide is applied outdoors to turf. Stay off the field until the flags are removed.
- if you are sensitive to chemicals, avoid the area of pesticide application for 72 hours.
- ensure that pets are kept away from the area of pesticide application

For indoor applications:

- cover or refrigerate edible products.
- remove or cover toys, clothes, and bedding from areas to be treated.
- remove pets including their food and water bowls and toys from the area to be treated.
- ventilate as much as possible during and, following an indoor pesticide application, open the windows.
- do not walk on treated areas and carpets until completely dry. Ask about drying times.

What types of pesticides will be applied?

Pesticide applicators may apply pesticides in several forms for control of insects and weeds. Dusts, aerosol sprays, sprays, baits, and fogs are all common forms in which pesticides exist and are used. For control of termites, the soil around the building may be impregnated with a pesticide. To control weeds, pesticides may be used as granules or sprays. Mechanical traps may also be used to control rodents.

In Massachusetts schools daycare centers and school age child care programs have to develop special pest management plans called Integrated Pest Management (IPM) plans. IPM is an approach to pest management which relies on a combination of common sense practices, including pesticides, for preventing and controlling pests. All plans are required to be submitted to the Department of Agricultural Resources. Check the MDAR website to see if your school has submitted its plan. <http://massnrc.org/ipm/index.html>

What if I have a question or problem?

Questions about what pesticides will be applied and why, and specific information about the application should be referred to the licensed pesticide applicator doing the work.

The Massachusetts Department of Agricultural Resources, Pesticide Enforcement is responsible for enforcing the pesticide regulations and laws. Contact Pesticide Enforcement at 617-626-1781. Additional information can be found at the Pesticide Programs website: <http://www.mass.gov/agr/pesticides/>

Updated August 2011

NPTN fact sheets are designed to answer questions that are commonly asked by the general public about pesticides that are regulated by the U.S. Environmental Protection Agency (US EPA). This document is intended to be educational in nature and helpful to consumers for making decisions about pesticide use.

National
Pesticide
Telecommunications
Network

Pyrethrins & Pyrethroids

The Pesticide Label: Labels provide directions for the proper use of a pesticide product. *Be sure to read the entire label before using any product.* A signal word on each product label indicates the product's short-term toxicity.

CAUTION- low toxicity

WARNING- moderate toxicity

DANGER- high toxicity

What are pyrethrins?

- Pyrethrins are insecticides that are derived from the extract of chrysanthemum flowers (pyrethrum) (1).
- The plant extract, called pyrethrum contains pyrethrin I and pyrethrin II collectively, called pyrethrins.
- Pyrethrins are widely used for control of various insect pests.

What are pyrethroids?

- Pyrethroids are synthetic (human-made) forms of pyrethrins. There are two types that differ in chemical structure and symptoms of exposure.
- Type I pyrethroids include allethrin, tetramethrin, resmethrin, d-phenothrin, bioresmethrin, and permethrin (1, 2).
- Some examples of type II pyrethroids are cypermethrin, cyfluthrin, deltamethrin, cyphenothrin, fenvalerate, and fluvalinate (1, 2).
- Both type I and II pyrethroids inhibit the nervous system of insects. This occurs at the sodium ion channels in the nerve cell membrane. Some type II pyrethroids also affect the action of a neurotransmitter called GABA (3).

How do pyrethrins (and pyrethroids) work?

- Nerve cell membranes have a specific electrical charge. Altering the amount of ions (charged atoms) passing through ion channels causes the membrane to depolarize which, in turn, causes a neurotransmitter to be released. Neurotransmitters help nerve cells communicate. Electrical messages sent between nerve cells allow them to generate a response, like a movement in an animal or insect.
- Pyrethrins affect the nervous system of insects by causing multiple action potentials in the nerve cells by delaying the closing of an ion channel (3).
- Pyrethrins and pyrethroids act as contact poisons, affecting the insect's nervous system (1, 4).
- Even though pyrethrins and pyrethroids are nerve poisons, they are not cholinesterase inhibitors like organophosphate or carbamate insecticides.
- Pesticide products containing pyrethrins usually contain a synergist (such as piperonyl butoxide). Synergists work by restricting an enzyme that insects use to detoxify the pyrethrins. A synergist allows the insecticide to be more effective (4).

There are many different types of pyrethroids, but the remainder of this fact sheet will deal with pyrethrins. Information on specific pyrethroids is available in other fact sheets.

What are some types of products that contain pyrethrins?

- indoor bugbombs or foggers
- human head-lice treatments
- pet flea sprays
- Dragon
- Drione
- Pyrenone
- Pyroicide

How toxic are pyrethrins?

Animals

- Pyrethrins are one of the least poisonous insecticides to mammals (2).
- Rats fed high doses (1,000 milligrams per kilogram of body weight or mg/kg) of pyrethrins showed liver damage (5).
- Rats exposed to pyrethrins exhibited difficulty or rapid breathing, incoordination, sprawling of limbs, tremors, aggression, sensitivity to external stimuli, twitching, and exhaustion (6). See box on **laboratory testing**.

Humans

- Inhaling pyrethrins can cause coughing, wheezing, shortness of breath, runny or stuffy nose, chest pain, or difficulty breathing (7).
- Skin contact can cause a rash, itching, or blisters (7).

Laboratory Testing: Before pesticides are registered by the U.S. EPA, they must undergo laboratory testing for short-term and long-term health effects. In these tests, laboratory animals are purposely fed a pesticide at high doses to cause toxic effects. These tests help scientists judge how these chemicals might affect humans, domestic animals, and wildlife in cases of overexposure. When pesticide products are used according to the label directions, toxic effects are not likely to occur because the amount of pesticide that people and animals may be exposed to is low compared to the doses fed to laboratory animals.

Effects of pyrethrins on human health and the environment depend on how much pyrethrins are present and the length and frequency of exposure. Effects also depend on the health of a person and/or certain environmental factors.

Do pyrethrins cause sensitization?

Animals

- The crude pyrethrum (initial plant extract) contains about 30 to 35 percent pyrethrins and about 50 percent impurities (2, 5).
- Various extracts from pyrethrum flowers have caused allergic contact dermatitis in sensitized and unsensitized guinea pigs (8). The commercially refined extract, which is present in insecticides today, did not produce any allergic reactions in guinea pigs (8, 9).
- Sensitization sometimes occurs in some individuals after a single exposure which causes either an asthmatic condition or a skin rash or inflammation. After the initial exposure to the sensitizing agent, the sensitized individual responds to a dose smaller than the initial dose.

Humans

- In one study, a person with a history of allergic contact dermatitis experimentally exposed to crude pyrethrum developed contact dermatitis, although this may have been caused by impurities in the extract (10).

Do pyrethrins break down and leave the body?

Animals

- Pyrethrins are low in toxicity to mammals because they are quickly broken down into inactive forms and pass from the body in the urine and feces (2, 5).

Humans

- Pyrethrum (the plant extract) may be absorbed by the digestive tract and the lungs. However, it is poorly absorbed through the skin (5).
- Based on animal studies, any amount of pyrethrins absorbed by humans would be expected to be rapidly excreted. Therefore, it is unlikely that pyrethrins

would accumulate in humans.

Are pyrethrins likely to cause cancer?

Animals

- In one study, rats were fed moderate to very high doses (100, 1000, or 3000 mg/kg) of pyrethrum (the plant extract) for 104 weeks. There was an increase in the non-cancerous (benign) thyroid tumors in females exposed to all doses and in males exposed to high to very high doses (11).
- In the same study, some females fed high doses (3000 mg/kg) of pyrethrum developed ovarian and benign liver tumors and males exposed to high doses (3000 mg/kg) developed benign parathyroid tumors and benign skin lesions.
- In another study, rats were fed low doses (up to 10 mg/kg) of pyrethrins, flavoring agents, and other pesticides showed no increase in tumors (6).

Humans

- Scientists have no data from work-related, accidental poisonings, or epidemiological studies that indicate whether or not pyrethrins are likely to cause cancer in humans.
- Initially, the Health Effects Division Carcinogenicity Peer Review Committee (CPRC) at the US EPA recently reviewed the carcinogenicity data of pyrethrins in animals and decided that they showed carcinogenicity. However, the CPRC could not classify pyrethrins into a carcinogenicity group until some of the tissue specimens from rats and mice were re-read. Subsequently, the CPRC will perform a second review of the carcinogenicity of pyrethrins (11).

Cancer: The U.S. EPA has strict guidelines that require testing of pesticides for their potential to cause cancer. These studies involve feeding laboratory animals large daily doses of the pesticide for up to 2 years. These animals are compared with a group of animals that did not receive the chemical. Animal studies help show whether a chemical is a potential human carcinogen. If a pesticide does not cause cancer in animal tests, then the EPA considers it unlikely the pesticide will cause cancer in humans.

Do pyrethrins cause reproductive problems or birth defects?

Animals

- Rabbits fed moderate doses (up to 90 mg/kg) of pyrethrins during a sensitive period of pregnancy had normal litters (5).
- Rats fed very high doses (5000 mg/kg) of pyrethrins for three weeks before their first mating produced low birth weight pups (5).
- There were no birth defects in pups of rabbits exposed to pyrethrins (12).

Humans

- There are no epidemiological, work-related or accidental exposure data on the potential of pyrethrins to cause reproductive problems or birth defects.

What happens to pyrethrins in the environment?

Soil

- Pyrethrins have a soil half-life of 12 days (13). They have an extremely low pesticide movement rating because they bind tightly to the soil (13). See box on **half-life**.

Photodegradation

- Pyrethrins are unstable in light or air (2). Pyrethrins are rapidly degraded in sunlight at the soil surface and in water.

Half-life is the time required for half of the compound to degrade.

1 half-life	=	50% degraded
2 half-lives	=	75% degraded
3 half-lives	=	88% degraded
4 half-lives	=	94% degraded
5 half-lives	=	97% degraded

Remember that the amount of chemical remaining after a half-life will always depend on the amount of the chemical originally applied.

What effects do pyrethrins have on wildlife?

- Pyrethrins are highly toxic to fish and tadpoles. They affect their skin touch receptors and balance organs (4).
- Pyrethrins are toxic to beneficial insect (such as honeybees) and many aquatic invertebrates (4).
- Pyrethrins are low in toxicity to humans, other mammals, and birds (4).

References

1. Klaassen, C. D., Amdur, M. O., & Doull, J. (Eds.). (1996). *Casarett & Doull's Toxicology. The Basic Science of Poisons* (5th ed.). Toronto: McGraw-Hill Companies, Inc.
2. Ray, D. E. (1991). Pesticides derived from plants and other organisms. In W. J. Hayes, Jr. & E. R. Laws (Eds.), *Handbook of Pesticide Toxicology. Vol. 2.* (pp.585-593). Toronto: Academic Press.
3. Costa, L. G. (1997). Basic Toxicology of Pesticides. In M. C. Keifer, M. D., M. P. H. (Ed.), *Human Health Effects of Pesticides. Occupational Medicine. State of the Art Reviews.* (Vol. 12 No. 2). (pp. 251-268). Philadelphia: Hanley & Belfus, Inc.
4. Tomlin, C. (Ed.). (1994). *A World Compendium. The Pesticide Manual. Incorporating the agrochemicals handbook.* (10th ed.). Bungay, Suffolk, U.K.: Crop Protection Publications.
5. Hayes, W. J. (1982). *Pesticides Studied in Man.* Baltimore: Williams & Wilkins.
6. Leahey, J. P. (Ed.). (1985). *The pyrethroid insecticides.* Philadelphia: Taylor & Francis Ltd.
7. Sittig, M. (1991). *Handbook of Toxic and Hazardous Chemicals and Carcinogens.* (3rd ed.). (Vol. 2). New Jersey: Noyes Publications.
8. Rickett, F. E., Tyszkiewicz, K., and Brown, N. C. (1972). *Pyrethrum dermatitis. I. The allergenic properties of various extracts of pyrethrum flowers.* Pestic. Sci. 3:57-66.
9. Rickett, F. E. and Tyszkiewicz, K. (1973). *Pyrethrum dermatitis. II. The allergenicity of pyrethrum oleoresin and its cross-reactions with saline extract of pyrethrum flowers.* Pestic. Sci. 4:801-810.
10. Mitchell, J. C., Dupuis, G. H. N. (1972). *Allergenic dermatitis from pyrethrum (Chrysanthemum spp.). The roles of pyrethrosin, a sesquiterpene lactone, and of pyrethrin II.* Br. J. Derm. 86:568-573.
11. United States Environmental Protection Agency (US EPA). Office of Prevention, Pesticides and Toxic Substances. *Carcinogenicity Peer Review of Pyrethrins.* February 22, 1995. Washington, DC.
12. Vettorazzi, G. (1979). *International Regulatory Aspects for Pesticide Chemicals. Toxicity Profiles.* (Vol.1). Boca Raton, Florida: CRC Press, Inc.
13. Wauchope, R. D., T. M. Butler, A. G. Hornsby, P. M.

Augustijn-Beckers, & J. P. Burt. (1992). The SCS/ARS/CES pesticide properties database for environmental decision making. [Online]. <ftp://ftp.nrcs.usda.gov/centers/itc/applications/wqmodels/gleams/>

For more information, call or write:
NPTN, Oregon State University, 333 Weniger
Hall, Corvallis, Oregon 97331-6502.
Phone: 1-800-858-7378
Fax: 1-541-737-0761
Email: nptn@ace.orst.edu
Internet: <http://ace.orst.edu/info/nptn/> or
<http://ace.orst.edu/info/extoxnet/>

Date reviewed: December 1998